

Drug Facts

Active ingredient

Brimonidine tartrate (0.025%).....Redness reliever

Purpose

Use

- relieves redness of the eye due to minor eye irritations

Warnings

For external use only

Do not use

- if solution changes color or becomes cloudy
- if single-use vial is not intact

Stop use and ask a doctor if

- you experience eye pain, changes in vision, continued redness or irritation of the eye
- condition worsens or persists for more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 5 years of age and over:
 - twist tab completely off to open vial
 - instill 1 drop in the affected eye(s)
 - discard vial immediately after use; do not re-use
 - store unused single-use vials in the child-resistant carton
 - do not use more than 4 times daily
 - remove contact lenses before use
 - wait at least 10 minutes before re-inserting contact lenses after use
 - if using other ophthalmic products while using this product, wait at least 5 minutes between each product
 - to avoid contamination, do not touch tip of container to any surface
- children under 5 years of age: consult a doctor

Other information

- store at 15-25 °C (59-77 °F)
- use only if pouch is sealed and single-use vial is intact
- this carton is child-resistant
- protect from light

Inactive ingredients

boric acid, calcium chloride dihydrate, glycerin, potassium chloride, sodium borate decahydrate, sodium chloride, water for injection. Hydrochloric acid and/or sodium hydroxide may be used to adjust pH.

Questions or comments? call: 1-800-553-5340